

# ArgiMent® AT

With Bimuno® Prebiotic

For Acute and Chronic Wounds

A Medical Food

Formulated for the nutritional management of pressure injuries and surgical wounds.

- 10 g Whey Protein
- 7 g Arginine & 7 g Glutamine
- 250 mg Vitamin C, 12.5 mg Zinc, 1 mg Copper & Prebiotic
- Nitrogen equivalent to 31.7 g of intact protein

## Orange Crème

Natural Flavor

Net Weight 1.5 oz (42.75 g)

Item 11220

Nutrition Facts	
Amount Per Serving	
Serving Size 1 Packet (42.75g)	
1 serving per container	
<b>Calories</b>	<b>120</b>
% Daily Value *	
Total Fat 0g	0%
Saturated Fat 0g	0%
Trans Fat 0g	
Cholesterol 25mg	8%
Sodium 40mg	2%
Potassium 87mg	2%
Phosphorus 35mg	2%
Total Carbohydrate 6g	2%
Dietary Fiber <1g	3%
Total Sugars 6g	
Includes 5g Added Sugars	10%
Protein 10g	20%
*The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice.	
† Daily Value not established.	



For clinical studies which support the use of this product as directed, visit [www.medtrition.com](http://www.medtrition.com)

info@medtrition.com  
www.medtrition.com

RH8B

Bimuno® is a registered trademark of Clasado IP Limited and is protected by patents 7,883,874, 8,058,047, 8,030,049 and 8,168,414. © 2018 Medtrition, Inc.

Distributed by

Medtrition, Inc.

P.O. Box 5387

Lancaster, PA 17606

United States

877.271.3570

**NOT FOR PARENTERAL USE**

Use under medical supervision.

renal function or liver disease.

at risk of dehydration, compromised

**CONTRAINDICATIONS:** Not for use by patients

Not a sole source of nutrition.

Billing Code: 94688-0112-20

HCPCS: B4155

Low in Lactose: 0.65 g/packet

Gluten free.

Glucanate, Yellow 6. **CONTAINS: MILK.**

Ascorbate, Sucralose, Zinc Sulfate, Copper

glucose, galactose), Orange Flavor, Calcium

(trans-galactooligosaccharide, lactose,

L-Arginine, L-Glutamine, Orange Juice

**INGREDIENTS:** Whey Protein Concentrate,

Powder, Fructose, Citric Acid, Bimuno®

**Do not mix with tube feeding formula.**

and after administration.

via syringe. Flush with 30 mL water before

Mixing with hot beverages may reduce potency of glutamine.

**Oral:** Mix 1 packet with 180–240 mL water or juice (two times daily).

**DOSING AND ADMINISTRATION:**

**ADVANCED THERAPY FOR ACUTE AND CHRONIC WOUNDS**