

# Banatotrol® plus

With Bimuno® Prebiotic

For Diarrhea and Loose Stools

## A Medical Food

Formulated for patients experiencing diarrhea and loose stools associated with the flu, antibiotics, tube feeding, oncology treatment or *C. diff*. Does not affect gut motility or cause constipation. Can be used as soon as *C. diff*. is suspected.

# Banana

Naturally Flavored

Net Weight 0.38 oz (10.75 g)  
Item 18470

**DOSING AND ADMINISTRATION:**  
**Oral: Diarrhea:** 1 packet of Banatotrol® Plus T1D (three times daily) added to beverages or foods, such as applesauce or yogurt. Maximum dose 6 packets/day with no more than 2 packets at a single time. Continue to use until diarrhea is resolved. Appropriate for continued long term use.  
**Loose Stools:** 1 packet 1-2 times per day as needed.

### Nutrition Facts

1 serving per container

Serving Size 1 Packet (10.75g)

Amount Per Serving

**40** Calories

% Daily Value\*

Total Fat 0g

Saturated Fat 0g

Trans Fat 0g

Cholesterol 0mg

Sodium 15mg

Potassium 125mg

Phosphorus 9mg

Total Carbohydrate 10g

Dietary Fiber 2g

Total Sugars 7g

Includes 1g Added Sugars

Protein 0g

Vit. D 0mcg 0%

Calcium 3mg 0%

Vit. K 0mg 0%

Iron 0mg 0%

\* The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice.



Distributed by  
Medtrition, Inc.  
P.O. Box 5387  
Lancaster, PA 17606  
United States  
877.271.3570  
info@medtrition.com  
www.medtrition.com

### NOT FOR PARENTERAL USE

Use under medical supervision.

**CONTRAINDICATIONS:** Latex Fruit Syndrome

Not a sole source of nutrition.

**HPCS:** B4104, Billing Code: 94688-0184-70

Low in Lactose: 0.27 g/packet

Gluten free.

**CONTAINS: MILK.**

(trans-galactooligosaccharides, lactose, glucose,

galactose).

**INGREDIENTS:** Banana Flakes, Bimuno®

When administering via feeding tube, we recommend

a size 1/4 French or larger to reduce the risk of clogging.

**Do not mix with tube feeding formula.**

Administer slowly via syringe. Flush with 30 mL water

before and after administration.

Banatotrol Plus T1D every 6 to 8 hours of feeding. Add

120 mL of room temperature water to a 240 mL cup.

Slowly stir in the contents of 1 packet. Mix well.

**Tube Feeding: G-Tube or J-Tube:** 1 packet of

For clinical studies which support the use of this product as directed, visit [www.medtrition.com](http://www.medtrition.com)

Bimuno® is a registered trademark of Clasado IP Limited and is protected by patents 7,883,874, 8,058,047, 8,030,049 and 8,168,414. ©2018 Medtrition, Inc.